

Compression Stockings



promoting
healthy skin

Champions for Skin Integrity



Replace compression stockings every 6 months or if they have a ladder or hole

Remove compression stockings immediately and seek advice if toes go purple or blue, the leg swells above or below the stockings, or you develop severe pain

If you remove compression stockings at night, reapply them first thing in the morning

Use a stocking applicator

Gently hand wash stockings, squeeze moisture out in a towel and dry in the shade

Wear rubber dishwashing gloves to help put your stockings on and to remove your stockings more easily



Do not wear rings, watches and jewellery when applying compression stockings

Do not leave any wrinkles in compression stockings





60 Musk Ave
Kelvin Grove Qld 4059
Brisbane, Australia

Email: ihbi@qut.edu.au

Email (Wound Healing): woundservice@qut.edu.au

www.ihbi.qut.edu.au

CRICOS No. 00213J

References:

RNAO, *Assessment and Management of Venous Leg Ulcers*, 2004, RNAO: Toronto

AWMA, *Australian and New Zealand Clinical Practice Guidelines for Prevention and Management of Venous Leg Ulcers*, 2011, AWMA: Barton. ACT

SIGN, *Management of chronic venous leg ulcers*, 2010, SIGN: Edinburgh

This project is funded by the Australian Government Department of Health and Ageing under the Encouraging Better Practice in Aged Care (EBPAC) program